Smith-Nephew

CORISurgical System

Hip Replacement Surgery

The CORI Surgical System is an advanced technology for hip replacement surgery. It provides detailed information for your unique hip anatomy, which can help your surgeon customize your replacement procedure. The results are improved patient outcomes¹⁻⁴ and higher patient satisfaction,⁴ when compared to conventional hip replacement.

your hip replacement surgery.



HALLIF FAME

Official joint replacement technology of the Pro Football Hall of Fame

Design of the CORI Surgical System

The CORI Surgical System supports computer-guided hip surgery. The system provides:

- Detailed data on your hip anatomy, including assessments of your bones
- Visuals to help determine the placement of your hip implant

The CORI Surgical System offers the ultimate combination: your surgeon's trained hand and the system's powerful technology.

Benefits of the CORI Surgical System

Your hip anatomy is unique – from the alignment to the measurement of the bones in your hip. The CORI Surgical System is designed to capture these details to help your surgeon optimize the placement of your hip implant.

The result is a personalized approach to surgery with significant benefits:

- Better accuracy in the surgical procedure, when compared to conventional (non-robotics) surgery⁵
- Improved patient outcomes, when compared to conventional (non-robotics)
 surgery¹⁻⁴
- Less chance of revision, when compared to conventional (non-robotics) surgery⁴
- Seamless integration with Smith+Nephew hip implants



To learn more about the CORI Surgical System, talk to your surgeon or visit **RediscoverYourGo.com**.



Not all patients are candidates for Smith+Nephew products. Hip replacement surgery is intended to relieve hip pain and improve hip function. However, implants may not produce the same feel or function as your original hip. There are potential risks with hip replacement surgery such as loosening, fracture, dislocation, wear and infection that may result in the need for additional surgery. Longevity of implants depends on many factors, such as types of activities and weight. Do not perform high impact activities such as running and jumping unless your surgeon tells you the bone has healed and these activities are acceptable. Early device failure, breakage or loosening may occur if you do not follow your surgeon's limitations on activity level. Early failure can happen if you do not guard your hip joint from overloading due to activity level, failure to control body weight, or accidents such as falls. Talk to your doctor to determine what treatment may be best for you. The information listed in this brochure is for informational purposes and is not meant as medical advice. For more information. please talk to your surgeon or visit RediscoverYourGo.com.

♦Trademark of Smith+Nephew. All Trademarks acknowledged. ©2023 Smith+Nephew. 39638 9/2023

1. Palit A, et al. Sci Rep. 2017;7:7238.
2. Weber M, et al. J Arthroplasty.
2016;31(11): 2514–2519.
3. Renkawitz T, et al. Bone Joint J.
2015;97:890–89.
4. Davis ET, et al. JBJS Open Access. 2021;6(3): e21.00006.
5. Naito Y, Hasegawa M, Tone S, Wakabayashi H, Sudo A. The accuracy of acetabular cup placement in primary total hip arthroplasty using an image-free navigation system. BMC Musculoskeletal Disorders. 2021;22(1).